

4  
626

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/19470</b> <b>(43) International Publication Date:</b> 22 April 1999 (22.04.99)
<b>(21) International Application Number:</b> PCT/US98/21444 <b>(22) International Filing Date:</b> 9 October 1998 (09.10.98)  <b>(30) Priority Data:</b> 08/948,276 9 October 1997 (09.10.97) US  <b>(71) Applicant:</b> THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 5th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US).  <b>(72) Inventor:</b> ERNST, Joel, D.; University of California, San Francisco, P.O. Box 0868, San Francisco, CA 94143-0868 (US).  <b>(74) Agent:</b> OSMAN, Richard, Aron; Science & Technology Law Group, 75 Denise Drive, Hillsborough, CA 94010 (US).		<b>(81) Designated States:</b> AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> GFP-ANNEXIN FUSION PROTEINS  <b>(57) Abstract</b>  Bifunctional green fluorescent protein (GFP)-annexin fusion proteins combine the inherent strong visible fluorescent properties of GFPs with the anionic phospholipid binding specificity of annexins. Recombinant host cells, especially bacteria, are used to efficiently express the fusion proteins in high yield and soluble form, suitable for rapid, one-step affinity purification. Uses include selective cellular and biochemical labeling, particularly anionic species, such as selectively labeling apoptotic cells.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

*GFP-Annexin Fusion Proteins*

The research carried out in the subject application was supported in part by NIH Grant # HL56001 (0577508-92-002). The government may have rights in any patent issuing on this application.

## INTRODUCTION

## Field of the Invention

The field of the invention is fluorescently-labeled proteins which specifically bind certain phospholipids.

## Background

The annexins are a family of proteins that specifically bind anionic phospholipids, including phosphatidylserine, in a calcium-dependent manner (Blackwood, R. A. and Ernst, J. D. (1990) *Biochem. J.* 266, 195-200; Seaton, B. A. (1995) *Annexins*, R. G. Landes, Austin, TX; *Cell Mol Life Sci*, Jun 1997;53(6), entire issue). While all annexins bind phosphatidylserine and calcium, they vary in their affinity for phosphatidylserine: for example, at saturating concentrations of calcium (i.e., = 1.0 mM), annexin V exhibits a 2- to 160-fold higher affinity for phosphatidylserine compared to other members of the annexin family (Tait, J. F., et al. (1988) *Biochemistry* 27, 6268-6276; Ernst, J. D., Mall, A. and Chew, G. (1994) *Biochem Biophys Res Com* 200, 867-876). Annexin binding specificities have been exploited for biological targeting (Tait et al., 1995, *J. Biol. Chem.* 270, 21594-21599; Oshawa et al, 1996, *J. Neurochem.* 67, 89-97; Okabayashi et al., 1996, *Gene* 177, 69-76).

Apoptosis, or programmed cell death, is a universal process that is important in development of multicellular organisms, regulation of the immune system, and clearance of abnormal (including neoplastic and virus-infected) cells (Thompson, C. B. (1995) *Science* 267, 1456-62). Among the early manifestations of apoptosis in all cell types studied to date is loss of the asymmetric distribution of plasma membrane phospholipids, which results in exposure of anionic phospholipids (including phosphatidylserine) on the extracellular leaflet of the plasma membrane. This exposure of phosphatidylserine, and thus apoptosis, can be detected by various methods, including binding of labeled annexins (Koopman, G., et al.

(1994) Blood 84, 1415-20; Martin, S. J., et al. (1995) J Exp Med 182, 1545-56; Broaddus, V. C., et al. (1996) J Clin Invest 98, 2050-2059; Zhang G, et al., 1997, Biotechniques, Sep;23(3):525-531. Recently, annexin binding specificity has been correlated with other cellular pathology, e.g. King KB (1997) J Cell Biochem 65(2), 131-144. Most studies to date have used FITC-annexin V and flow cytometry to identify and enumerate apoptotic cells. Labeling annexin V with FITC requires multiple manipulations of the protein and results in a heterogeneous mixture of labeled protein molecules which vary in the number and position of bound FITC molecules. Moreover, the amino acid residue of annexin V that is most readily available for labeling by FITC is on or near the phospholipid-binding surface, which results in quenching of FITC-annexin V fluorescence by 40-50% upon binding phospholipid membranes (Tait, 1988; Ernst, 1994; supra).

In an effort to circumvent these limitations of FITC-annexins, the present inventor sought to prepare annexins that were labeled homogeneously and that did not change fluorescence properties upon binding membrane phospholipids. Described herein are the preparation and characterization of endogenously fluorescent phosphatidylserine-binding proteins containing *Aequorea victoria* green fluorescent proteins (GFPs) fused to annexins. It is shown that these reagents offer highly sensitive detection of apoptotic cells by flow cytometry or fluorescent microscopy, and offer several advantages to chemically modified annexins.

## SUMMARY OF THE INVENTION

The invention provides methods and compositions relating to *Aequorea victoria* GFP-annexin fusion proteins; particularly, recombinant polypeptides comprising a bifunctional green fluorescent protein - annexin fusion protein providing an equivalent or enhanced measured fluorescent property of the green fluorescent protein and an equivalent or enhanced measured binding specificity of the annexin. In a particular embodiment, the fusion protein comprises a full-length N-terminal GFP fused to a full-length annexin V through a linker comprising an alanine, wherein the fused GFP and annexin moieties provide greater or equal fluorescent intensity and anionic phospholipid binding affinity, respectively, than do the corresponding unfused GFP and annexin proteins.

The invention also provides host cells expressing the subject proteins, including bacteria expressing the subject proteins in soluble form, and methods of using such cells to

make the fusion proteins. Uses of the subject fusion proteins include selective cellular and biochemical labeling, particularly anionic species, such as anionic phospholipids. In a particular embodiment, the fusion proteins are used to selectively label apoptotic, dead and/or injured cells.

## DESCRIPTION OF PARTICULAR EMBODIMENTS OF THE INVENTION

The subject bifunctional GFP-annexin fusion proteins combine the inherent intense fluorescent properties of green fluorescent proteins with the binding specificity of annexins. The GFPs derive from the jellyfish *Aequorea victoria*; see e.g. US Patent No.5,491,084 for definition, and include variants offering a variety of different excitation and emission wavelengths; see e.g. Heim and Tsein, 1996, Current Biology 6, 178-182. The GFP moiety of the fusion proteins provide an equivalent or enhanced measured qualitative and/or quantitative fluorescent property compared with the corresponding unfused GFP protein. Preferred fluorescent properties are emission and/or excitation peaks, preferably an maximum fluorescent emission peak in unchanged or detectably optimized wavelength and/or undiminished or enhanced in magnitude or total intensity.

The subject annexins may be derived from a variety of eukaryotic sources, see e.g. Cell Mol Life Sci, June 1997;53(6), entire issue, esp. Liemann S, Huber R, at 516-521 and Morgan RO, at 508-515, and any of the at least thirteen distinct annexin types may be used. The annexin moiety of the fusion proteins provide an equivalent or enhanced measured qualitative and/or quantitative binding specificity compared with the corresponding unfused annexin protein. Preferred binding specificities have equivalent or enhanced affinity for particular anionic cellular components, particularly phospholipids, such as phosphatidylserine.

The GFP and annexin moieties may be separated by a linker peptide, typically from about 1 to 50 residues, which facilitates or at least does not interfere with the requisite bifunctionality of the fusion proteins. The linker may enhance the conformational opportunities of the GFP and annexin moieties and/or provide a third functionality to the fusion protein, e.g. epitopes, post-translational processing sites, etc.. Exemplary linkers include alanine or polyalanine, glycine or polyglycine, epitope tags such as FLAG, processing sites such as phosphorylation, ubiquitination or protease recognition/cleavage sites, etc.

Exemplary bifunctional fusion proteins are shown in Table I.

Ref No.	GFP Variant Moiety	Annexin Moiety, residues	Linker	Activity	
				GFP	Annexin
5	T65V-I	S65T	hV (SEQ ID NO:1, residues 1-320)	ϕ	+++
	T65V-II	S65T	hV (SEQ ID NO:1, residues 12-320)	Gly	+++
	T65V-III	S65T	hV (SEQ ID NO:1, residues 3-319)	Ala	+++
	T65V-IV	S65T	mV (SEQ ID NO:2, residues 1-319)	(Ala) <sub>3</sub>	+++
	T65V-V	S65T	rV (SEQ ID NO:3, residues 12-319)	(Ala) <sub>9</sub>	+++
10	T65IV-I	S65T	hIV (SEQ ID NO:4, residues 1-321)	FLAG	+++
	T65III-I	S65T	hIII (SEQ ID NO:5, residues 1-323)	ϕ	+++
	H66V-I	Y66H	hV (SEQ ID NO:1, residues 12-320)	Gly	+++
	W66V-I	Y66W	hV (SEQ ID NO:1, residues 6-320)	(AlaGly) <sub>2</sub>	+++
	L64III-I	F64L	hIII (SEQ ID NO:1, residues 6-323)	Ala	+++
15	H66I-I	S65T	hI (SEQ ID NO:6, residues 1-346)	GlyAlaGly	+++
	T65I-I	S65T	hI (SEQ ID NO:6, residues 41-346)	AlaSerAla	+++

The invention provides recombinant nucleic acids encoding the subject fusion proteins. Typically, natural isolated nucleic acids encoding the GFP and annexin moieties are spliced into expression constructs using conventional methodologies, see e.g. Molecular Cloning, A Laboratory Manual (Sambrook, *et al.* Cold Spring Harbor Laboratory), Current Protocols in Molecular Biology (Eds. Ausubel, *et al.*, Greene Publ. Assoc., Wiley-Interscience, NY) and references cited herein. Alternatively, the amino acid sequences of the subject peptides are used to back-translate peptide-encoding nucleic acids optimized for selected expression systems (Holler et al. (1993) Gene 136, 323-328; Martin et al. (1995) Gene 154, 150-166). In either instance, the constructs are designed for expression in any conventional system, such as bacterial, insect, plant and mammalian expression systems. The proteins are preferably secreted and/or expressed in soluble form; preferably most of the protein secreted by or retained within the host cell is in soluble form. Preferred soluble expression avoids denaturation/renaturation and permits single step affinity purification of >90%, preferably >95%, preferably in a yield of at least 10, more preferably at least 25 mg/L. In a particular embodiment, the temperature of the expressing host is reduced at least

5, preferably at least 10, more preferably at least 15°C below physiological or environmental temperature for the host (e.g. below 37°C for E.coli or human cells).

Uses of the subject fusion proteins include selective cellular and biochemical labeling, particularly anionic species, such as anionic phospholipids. The subject proteins may be exposed to the targeted cellular or biochemical component in any convenient way, e.g. direct exogenous addition, indirectly by introduction into a cell and expression of a fusion protein encoding nucleic acid, etc., In a particular embodiment, the fusion proteins are used to selectively label apoptotic, dead and/or injured cells.

Without further description, one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure. Other generic configurations will be apparent to one skilled in the art. All publications and patent applications cited in this specification, and cited references therein, are herein incorporated by reference as if each individual publication, patent application or reference were specifically and individually indicated to be incorporated by reference.

### EXAMPLES

*Plasmid construction.* Parental annexin plasmids for construction of GFP annexin fusions were constructed as previously described in published literature (e.g. Seaton, 1995, supra and citations therein). For example, the parent plasmid for construction of the GFP-annexin V fusion was pET9dE2, which were previously constructed and used for expression of human annexin V (Ernst, 1994, supra). Similarly, constructs encoding a panel of GFPs are constructed using commercially available and/or published material and methods (e.g. US Patent No. 5,491,084; Clontech, Apr 1997; Kaln, 1997, Biotechnol. International 8/97; Heim and Tsien, 1996, Current Biology 6, 178-182). For example, the open reading frame of green fluorescent protein (S65T variant) was amplified by PCR using pS65T-C1 (Clontech) as template and primers designed to incorporate sites for Rca I for ligation to the Nco I site of pET9dE2. In this case, the forward primer introduces a change in codon 2 to encode serine (as in wild-type GFP) rather than the glycine encoded by the template plasmid; and the backward primer eliminates the stop codon at the 3' end of the GFP open reading

frame and introduces an alanine codon to form the junction with the 5' end of annexin V. After digestion of the PCR product with Rca I, it was ligated to Nco I-digested pET9dE2 and used to transform *E. coli* DH5 $\alpha$ , and transformants that contained the GFP fragment in the correct orientation were used to transform *E. coli* BL-21(DE3) for protein expression.

5        *Expression and purification of protein.* Initial attempts at protein expression under standard conditions (growth at 37°C, induction with 0.4 mM IPTG) yielded brightly fluorescent but insoluble protein. Consequently, procedures were devised for expression of soluble, bifunctional proteins. In one embodiment, the GFP-annexin fusion proteins were expressed in *E. coli* BL-21(DE3) by growth in LB media at room temperature for 18-20  
10        hours, without IPTG induction. This resulted in fusion proteins that were bright green and present in the soluble fraction (approximately 80%) after lysozyme digestion and probe sonication of the bacterial suspension. After adding calcium (2.5 mM) to the supernatant fraction of the *E. coli* lysate, GFP-annexins were purified by affinity chromatography using phospholipids (e.g. phosphatidylserine) immobilized on controlled-pore glass (Ernst, J. D.,  
15        et al., (1991) J. Biol. Chem. 266, 6670-6673; Ernst, J. D. (1991) J. Immunol. 146, 3110-3114; Ernst, 1994).

*Characterization of GFP-annexin.* Fluorescence excitation and emission spectra were obtained using an SLM8000C spectrofluorimeter equipped with motorized excitation and emission monochromators and bandpass settings of 4 nm. Phospholipid-containing  
20        liposomes were prepared as previously described (Ernst, 1994, supra).

*Studies of apoptotic cells.* Rabbit and human pleural mesothelial cells were plated in 6 well plates at near confluency. For the experiments detailed below, cells were incubated overnight in the apoptotic stimuli, crocidolite (5-20  $\mu\text{g}/\text{cm}^2$ ) or actinomycin D (0.33  $\mu\text{M}$ ), in serum-free conditions to avoid serum coating onto the asbestos fiber. After floating cells  
25        were collected by aspiration of medium, attached cells were detached with trypsin (2.5%) and added to the floating cells. Cells were kept on ice to minimize ongoing apoptosis. Cells exposed to asbestos fibers were filtered through a 100  $\mu\text{m}$  cell strainer to remove fibers prior to flow cytometric analysis. Cells (approximately  $2.5 \times 10^5$  cells per condition) were then  
30        spun and resuspended in 200  $\mu\text{l}$  of annexin buffer (Hank's buffer with 15 mM Hepes and 2 mM total calcium concentration [1.3 mM in the Hank's plus an additional 0.7 mM  $\text{CaCl}_2$ ]).

      Cells ( $2.5 \times 10^5$  in 200  $\mu\text{l}$ ) were incubated with GFP-annexin or FITC-annexin (both at 3  $\mu\text{g}/\text{ml}$ ) for 10 min on ice. Immediately prior to analysis by flow cytometry, propidium



iodide (15  $\mu$ g/ml, Sigma Chemical Co.) was added to each tube. No further washing and no fixation was performed.

Cells on ice were analysed by flow cytometry (FACSort, Becton Dickinson, San Jose, CA) with acquisition and data analysis using CELLQuest Software (Becton Dickinson). Compensation for the use of two fluorescent probes was set using control cells stained with either GFP-annexin or propidium iodide alone. 10,000 events per sample were acquired to ensure adequate mean data.

For detection of intracellular antigens, cells require fixation and permeabilization. To determine if GFP-annexin binding would be altered by those conditions, various fixatives and permeabilization were tested. During these studies, the calcium concentration was increased to 5 mM. After cells were stained with GFP-annexin as above, cells were washed twice in annexin buffer to remove all free GFP-annexin. To determine the stability of the bound GFP-annexin during fixation, the cells were then fixed with either glutaraldehyde (0.5-2%) or paraformaldehyde (2-4 %) for 10 min in the dark. The cells were then washed in annexin buffer (with 5 mM  $\text{CaCl}_2$ ) and permeabilized with Triton 0.1% in the same buffer for 4 min. Following an additional wash, cells were resuspended in annexin buffer (with 5 mM  $\text{CaCl}_2$ ) for analysis of GFP-annexin binding.

*Expression and purification of GFP-annexins:* Under the low temperature expression conditions, approximately 80% of the GFP-annexins expressed were present in the soluble fraction of the *E. coli* lysate, and generally >90% could be isolated to purity in single step phospholipid affinity chromatography. For example, GFP-annexin V proteins were isolated to >90% purity in a single step by calcium-dependent phosphatidylserine affinity chromatography. This served as an efficient purification step as well as providing evidence that the phospholipid binding ability of the annexin domains were preserved in these chimera. Another chimeric protein in which GFP was fused to the carboxyl terminus of annexin V was also soluble and fluorescent, but did not bind phosphatidylserine. The isolated proteins had the electrophoretic mobility predicted by amino acid compositions (61 kDa for annexin V fusions), were recognized by an antibody to the annexin moiety, and exhibited SDS-resistant fluorescence with UV transillumination of the gel and exhibited the functional properties of both antecedents. The failure of an annexin V fusion protein that contained the moieties in the contrasting orientation (e.g., GFP fused to the carboxyl terminus of annexin V) to bind phospholipids was unexpected and may have been the result

of the GFP domain masking the phospholipid binding surface of the annexin.

*Fluorescence properties of GFP-annexins:* To assure that the fusion of annexins to GFPs did not reduce the fluorescence properties of the GFP moieties, the excitation and emission spectra of the GFP-annexin fusion proteins were examined and the spectra of the GFPs found undiminished. For example, the GFP(S65T)-annexin V fusion proteins exhibited a discrete peak of fluorescence emission centered at 512 nm when excited at 490 nm. Consistent with the prior observation that the alteration of residue 65 from serine to threonine causes loss of excitation of GFP in the near ultraviolet range, there was no fluorescence detected when these GFP-annexin V chimeras were excited at 375 nm. When fluorescence emission was monitored at 510 nm, there was a single major excitation peak at 465-495 nm, with a minor trough at 480 nm. The fluorescence spectra were unaffected by the addition of calcium (1.0 mM), phosphatidylserine-containing liposomes (1.0  $\mu$ M), or both. These GFP-containing proteins are ideally suited for use with the argon-laser based flow cytometers (excitation at 488 nm/detection at 530 nm).

*Use of GFP-annexin V to detect apoptotic cells by flow cytometry.* To determine whether the annexin V chimeras retained the ability of annexin V to bind apoptotic cells in a specific and calcium-dependent manner, a well-characterized model system of apoptosis of pleural mesothelial cells was used (Broadbush, 1996, supra). GFP-annexin V chimeras did not bind to normal cells, but exhibited saturable binding to apoptotic mesothelial cells as detected by flow cytometry. Compared to FITC-annexin V, fluorescence of GFP-annexin V chimera-labeled apoptotic cells was 5 times brighter, despite using GFP-annexin V chimeras at approximately 1/2 the molar concentration of FITC-annexin V. In these experiments, staining with either FITC-annexin V or GFP-annexin V chimeras provided a more sensitive detection of apoptotic cells than staining with acridine orange. Binding of GFP-annexin V chimeras to apoptotic cells was calcium-dependent, and could be competitively antagonized by unlabeled annexin V. Therefore, in addition to retaining the fluorescence properties of GFP, GFP-annexin V chimeras also retain the ability of annexin V to specifically detect exposure of anionic phospholipids on the outer leaflet of the plasma membrane of apoptotic cells. GFP-annexin V chimeras could also be used in experiments in which exclusion of propidium iodide (PI) was used to distinguish early apoptotic (GFP-annexin V+, PI-) cells from late apoptotic or necrotic cells (GFP-annexin V+, PI+).

In studies of apoptosis in vivo, it may be important to identify the cell type

undergoing apoptosis. One means of phenotyping apoptosis cells is to costain with an antibody that identifies a specific cell type. Some cells, for example mesothelial cells have no unique cell surface antigen and must be identified by intracellular expression of intermediate filaments, namely cytokeratin. Access of antibodies to this or other intracellular antigens requires cell permeabilization with detergent, which disrupts the integrity of plasma membrane phospholipids. When apoptotic cells were first labeled with GFP-annexin V chimeras, washed to remove unbound fusion protein, fixed with paraformaldehyde or glutaraldehyde, and permeabilized with 0.1% Triton X-100, GFP-annexin V chimera binding was retained. Moreover, after staining with a monoclonal antibody to cytokeratin and PE-labeled goat anti-mouse IgG, two-color flow cytometry could be used to identify and enumerate apoptotic mesothelial cells in a mixture of cells (lymphocytes, macrophages, and neutrophils) obtained from the pleural space of rabbits treated with crocidolite asbestos.

Due to their bifunctional properties, GFP-annexins are useful reagents for further studies of apoptosis and of disorders of the erythrocyte membrane that are characterized by loss of phospholipid asymmetry and exposure of anionic phospholipids (Kuypers, F. A., et al. (1996) *Blood* 87, 1179-1187; Wood, B. L., Gibson, D. F. and Tait, J. F. (1996) *Blood* 88, 1873-80). In addition to the utility of GFP-annexins as reagents for studying membrane phospholipids, the approach described here is useful in constructing annexin fusions to GFP to study the interaction of specific annexins with intracellular membranes in intact cells. Annexins have been found to interact with membranes of phagosomes, endosomes, and intracellular vesicles containing microbial pathogens (Ernst, 1991, *supra*; Emans, N., et al. (1993) *J Cell Biol* 120, 1357-1369; Desjardins, M., et al. (1994) *J Biol Chem* 269, 32194-200; Majeed, M., et al. (1994) *Infect Immun* 62, 127-134; Diakonova, M., et al. (1997) *J Cell Sci* 110, 1199-213). The ability to use bifunctional GFP-annexin fusion proteins that are functionally well-characterized to study the interaction of annexins with intracellular membranes will further advance the understanding of the intracellular functions of the annexin proteins.

## WHAT IS CLAIMED IS:

1. A recombinant polypeptide comprising a bifunctional *Aequorea victoria* green fluorescent protein - annexin fusion protein, said fusion protein comprising GFP and annexin moieties which provide greater or equal fluorescent intensity and anionic phospholipid binding affinity, respectively, than do the corresponding unfused GFP and annexin proteins.

2. The polypeptide of claim 1, wherein the GFP and annexin moieties are selected from: S65T GFP variant/hAnnexinV, residues 1-320 (SEQ ID NO:1, residues 1-320); S65T GFP variant/hAnnexinV, residues 12-320 (SEQ ID NO:1, residues 12-320); S65T GFP variant/hAnnexinV, residues 3-319 (SEQ ID NO:1, residues 3-319); S65T GFP variant/mAnnexinV, residues 1-319 (SEQ ID NO:2, residues 1-319); S65T GFP variant/rAnnexinV, residues 12-319 (SEQ ID NO:3, residues 1-319); S65T GFP variant/hAnnexinIV, residues 1-321 (SEQ ID NO:4, residues 1-321); S65T GFP variant/hAnnexinIII, residues 1-323 (SEQ ID NO:5, residues 1-323); Y66H GFP variant/hAnnexinV, residues 12-320 (SEQ ID NO:1, residues 12-320); Y66W GFP variant/hAnnexinV, residues 6-320 (SEQ ID NO:1, residues 6-320); F64L GFP variant/hAnnexinIII, residues 6-323 (SEQ ID NO:5, residues 6-323); S65T GFP variant/hAnnexinI, residues 1-346 (SEQ ID NO:6, residues 1-346); and S65T GFP variant/hAnnexinI, residues 41-346 (SEQ ID NO:6, residues 41-346).

3. A bacterium comprising a soluble polypeptide according to claim 1.

4. A method of making a polypeptide comprising a bifunctional green fluorescent protein - annexin fusion protein, said method comprising the steps of culturing a bacterium comprising a nucleic acid encoding a polypeptide according to claim 1 under conditions wherein the polypeptide is solubly expressed within the bacterium.

5. A method of labeling a cell comprising the step of contacting the cell with a polypeptide according to claim 1.

6. A cell which expresses the polypeptide of claim 1.

7. A mammalian cell which expresses the polypeptide of claim 1.
8. A cultured cell which expresses the polypeptide of claim 1.

## SEQUENCE LISTING

&lt;110&gt; Ernst, Joel D.

&lt;120&gt; GFP-Annexin Fusion Proteins

&lt;130&gt; UCSF97-113

&lt;140&gt; 08/948,276

&lt;141&gt; 1997-10-09

&lt;160&gt; 6

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 320

&lt;212&gt; PRT

&lt;213&gt; human

&lt;400&gt; 1

Met Ala Gln Val Leu Arg Gly Thr Val Thr Asp Phe Pro Gly Phe Asp  
 1 5 10 15

Glu Arg Ala Asp Ala Glu Thr Leu Arg Lys Ala Met Lys Gly Leu Gly  
 20 25 30

Thr Asp Glu Glu Ser Ile Leu Thr Leu Leu Thr Ser Arg Ser Asn Ala  
 35 40 45

Gln Arg Gln Glu Ile Ser Ala Ala Phe Lys Thr Leu Phe Gly Arg Asp  
 50 55 60

Leu Leu Asp Asp Leu Lys Ser Glu Leu Thr Gly Lys Phe Glu Lys Leu  
 65 70 75 80

Ile Val Ala Leu Met Lys Pro Ser Arg Leu Tyr Asp Ala Tyr Glu Leu  
 85 90 95

Lys His Ala Leu Lys Gly Ala Gly Thr Asn Glu Lys Val Leu Thr Glu  
 100 105 110

Ile Ile Ala Ser Arg Thr Pro Glu Glu Leu Arg Ala Ile Lys Gln Val  
 115 120 125

Tyr Glu Glu Glu Tyr Gly Ser Ser Leu Glu Asp Asp Val Val Gly Asp  
 130 135 140

Thr Ser Gly Tyr Tyr Gln Arg Met Leu Val Val Leu Leu Gln Ala Asn  
 145 150 155 160  
 5 Arg Asp Pro Asp Ala Gly Ile Asp Glu Ala Gln Val Glu Gln Asp Ala  
 165 170 175  
 Gln Ala Leu Phe Gln Ala Gly Glu Leu Lys Trp Gly Thr Asp Glu Glu  
 180 185 190  
 10 Lys Phe Ile Thr Ile Phe Gly Thr Arg Ser Val Ser His Leu Arg Lys  
 195 200 205  
 Val Phe Asp Lys Tyr Met Thr Ile Ser Gly Phe Gln Ile Glu Glu Thr  
 15 210 215 220  
 Ile Asp Arg Glu Thr Ser Gly Asn Leu Glu Gln Leu Leu Leu Ala Val  
 225 230 235 240  
 20 Val Lys Ser Ile Arg Ser Ile Pro Ala Tyr Leu Ala Glu Thr Leu Tyr  
 245 250 255  
 Tyr Ala Met Lys Gly Ala Gly Thr Asp Asp His Thr Leu Ile Arg Val  
 260 265 270  
 25 Met Val Ser Arg Ser Glu Ile Asp Leu Phe Asn Ile Arg Lys Glu Phe  
 275 280 285  
 Arg Lys Asn Phe Ala Thr Ser Leu Tyr Ser Met Ile Lys Gly Asp Thr  
 30 290 295 300  
 Ser Gly Asp Tyr Lys Lys Ala Leu Leu Leu Leu Cys Gly Glu Asp Asp  
 305 310 315 320  
 35 <210> 2  
 <211> 319  
 <212> PRT  
 <213> mouse  
 <400> 2  
 40 Met Ala Thr Arg Gly Thr Val Thr Asp Phe Pro Gly Phe Asp Gly Arg  
 1 5 10 15

	Ala Asp Ala Glu Val Leu Arg Lys Ala Met Lys Gly Leu Gly Thr Asp	
	20	25 30
5	Glu Asp Ser Ile Leu Asn Leu Leu Thr Ser Arg Ser Asn Ala Gln Arg	
	35	40 45
	Gln Glu Ile Ala Gln Glu Phe Lys Thr Leu Phe Gly Arg Asp Leu Val	
	50	55 60
10	Asp Asp Leu Lys Ser Glu Leu Thr Gly Lys Phe Glu Lys Leu Ile Val	
	65	70 75 80
	Ala Met Met Lys Pro Ser Arg Leu Tyr Asp Ala Tyr Glu Leu Lys His	
	85	90 95
15	Ala Leu Lys Gly Ala Gly Thr Asp Glu Lys Val Leu Thr Glu Ile Ile	
	100	105 110
	Ala Ser Arg Thr Pro Glu Glu Leu Ser Ala Ile Lys Gln Val Tyr Glu	
20	115	120 125
	Glu Glu Tyr Gly Ser Asn Leu Glu Asp Asp Val Val Gly Asp Thr Ser	
	130	135 140
25	Gly Tyr Tyr Gln Arg Met Leu Val Val Leu Leu Gln Ala Asn Arg Asp	
	145	150 155 160
	Pro Asp Thr Ala Ile Asp Asp Ala Gln Val Glu Leu Asp Ala Gln Ala	
	165	170 175
30	Leu Phe Gln Ala Gly Glu Leu Lys Trp Gly Thr Asp Glu Glu Lys Phe	
	180	185 190
	Ile Thr Ile Phe Gly Thr Arg Ser Val Ser His Leu Arg Arg Val Phe	
35	195	200 205
	Asp Lys Tyr Met Thr Ile Ser Gly Phe Gln Ile Glu Glu Thr Ile Asp	
	210	215 220
40	Arg Glu Thr Ser Gly Asn Leu Glu Gln Leu Leu Leu Ala Val Val Lys	
	225	230 235 240



Ser Ile Arg Ser Ile Pro Ala Tyr Leu Ala Glu Thr Leu Tyr Tyr Ala  
 245 250 255  
 Met Lys Gly Ala Gly Thr Asp Asp His Thr Leu Ile Arg Val Val Val  
 5 260 265 270  
 Ser Arg Ser Glu Ile Asp Leu Phe Asn Ile Arg Lys Glu Phe Arg Lys  
 275 280 285  
 10 Asn Phe Ala Thr Ser Leu Tyr Ser Met Ile Lys Gly Asp Thr Ser Gly  
 290 295 300  
 Asp Tyr Lys Lys Ala Leu Leu Leu Leu Cys Gly Gly Glu Asp Asp  
 15 305 310 315  
 <210> 3  
 <211> 318  
 <212> PRT  
 <213> rat  
 20 <400> 3  
 Ala Leu Arg Gly Thr Val Thr Asp Phe Ser Gly Phe Asp Gly Arg Ala  
 1 5 10 15  
 Asp Ala Glu Val Leu Arg Lys Ala Met Lys Gly Leu Gly Thr Asp Glu  
 25 20 25 30  
 Asp Ser Ile Leu Asn Leu Leu Thr Ala Arg Ser Asn Ala Gln Arg Gln  
 35 40 45  
 30 Gln Ile Ala Glu Glu Phe Lys Thr Leu Phe Gly Arg Asp Leu Val Asn  
 50 55 60  
 Asp Met Lys Ser Glu Leu Thr Gly Lys Phe Glu Lys Leu Ile Val Ala  
 65 70 75 80  
 35 Leu Met Lys Pro Ser Arg Leu Tyr Asp Ala Tyr Glu Leu Lys His Ala  
 85 90 95  
 Leu Lys Gly Ala Gly Thr Asp Glu Lys Val Leu Thr Glu Ile Ile Ala  
 40 100 105 110  
 Ser Arg Thr Pro Glu Glu Leu Arg Ala Ile Lys Gln Ala Tyr Glu Glu

	115	120	125
	Glu Tyr Gly Ser Asn Leu	Glu Asp Asp Val Val	Gly Asp Thr Ser Gly
	130	135	140
5	Tyr Tyr Gln Arg Met Leu Val Val Leu Leu Gln Ala Asn Arg Asp Pro		
	145	150	155 160
	Asp Thr Ala Ile Asp Asp Ala Gln Val Glu Leu Asp Ala Gln Ala Leu		
10	165	170	175
	Phe Gln Ala Gly Glu Leu Lys Trp Gly Thr Asp Glu Glu Lys Phe Ile		
	180	185	190
15	Thr Ile Leu Gly Thr Arg Ser Val Ser His Leu Arg Arg Val Phe Asp		
	195	200	205
	Lys Tyr Met Thr Ile Ser Gly Phe Gln Ile Glu Glu Thr Ile Asp Arg		
	210	215	220
20	Glu Thr Ser Gly Asn Leu Glu Asn Leu Leu Leu Ala Val Val Lys Ser		
	225	230	235 240
	Ile Arg Ser Ile Pro Ala Tyr Leu Ala Glu Thr Leu Tyr Tyr Ala Met		
25	245	250	255
	Lys Gly Ala Gly Thr Asp Asp His Thr Leu Ile Arg Val Ile Val Ser		
	260	265	270
30	Arg Ser Glu Ile Asp Leu Phe Asn Ile Arg Lys Glu Phe Arg Lys Asn		
	275	280	285
	Phe Ala Thr Ser Leu Tyr Ser Met Ile Lys Gly Asp Thr Ser Gly Asp		
	290	295	300
35	Tyr Lys Lys Ala Leu Leu Leu Leu Cys Gly Gly Glu Asp Asp		
	305	310	315
	<210> 4		
40	<211> 321		
	<212> PRT		
	<213> human		

&lt;400&gt; 4

Met Ala Met Ala Thr Lys Gly Gly Thr Val Lys Ala Ala Ser Gly Phe  
 1 5 10 15

5 Asn Ala Met Glu Asp Ala Gln Thr Leu Arg Lys Ala Met Lys Gly Leu  
 20 25 30

Gly Thr Asp Glu Asp Ala Ile Ile Ser Val Leu Ala Tyr Arg Asn Thr  
 35 40 45

10 Ala Gln Arg Gln Glu Ile Arg Thr Ala Tyr Lys Ser Thr Ile Gly Arg  
 50 55 60

15 Asp Leu Ile Asp Asp Leu Lys Ser Glu Leu Ser Gly Asn Phe Glu Gln  
 65 70 75 80

Val Ile Val Gly Met Met Thr Pro Thr Val Leu Tyr Asp Val Gln Glu  
 85 90 95

20 Leu Arg Arg Ala Met Lys Gly Ala Gly Thr Asp Glu Gly Cys Leu Ile  
 100 105 110

Glu Ile Leu Ala Ser Arg Thr Pro Glu Glu Ile Arg Arg Ile Ser Gln  
 115 120 125

25 Thr Tyr Gln Gln Gln Tyr Gly Arg Ser Leu Glu Asp Asp Ile Arg Ser  
 130 135 140

30 Asp Thr Ser Phe Met Phe Gln Arg Val Leu Val Ser Leu Ser Ala Gly  
 145 150 155 160

Gly Arg Asp Glu Gly Asn Tyr Leu Asp Asp Ala Leu Val Arg Gln Asp  
 165 170 175

35 Ala Gln Asp Leu Tyr Glu Ala Gly Glu Lys Lys Trp Gly Thr Asp Glu  
 180 185 190

Val Lys Phe Leu Thr Val Leu Cys Ser Arg Asn Arg Asn His Leu Leu  
 195 200 205

40 His Val Phe Asp Glu Tyr Lys Arg Ile Ser Gln Lys Asp Ile Glu Gln  
 210 215 220

Ser Ile Lys Ser Glu Thr Ser Gly Ser Phe Glu Asp Ala Leu Leu Ala  
 225 230 235 240  
 5 Ile Val Lys Cys Met Arg Asn Lys Ser Ala Tyr Phe Ala Glu Lys Leu  
 245 250 255  
 Tyr Lys Ser Met Lys Gly Leu Gly Thr Asp Asp Asn Thr Leu Ile Arg  
 260 265 270  
 10 Val Met Val Ser Arg Ala Glu Ile Asp Met Leu Asp Ile Arg Ala His  
 275 280 285  
 Phe Lys Arg Leu Tyr Gly Lys Ser Leu Tyr Ser Phe Ile Lys Gly Asp  
 15 290 295 300  
 Thr Ser Gly Asp Tyr Arg Lys Val Leu Leu Val Leu Cys Gly Gly Asp  
 305 310 315 320  
 20 Asp  
 <210> 5  
 <211> 323  
 <212> PRT  
 25 <213> human  
 <400> 5  
 Ser Ala Ser Ile Trp Val Gly His Arg Gly Thr Val Arg Asp Tyr Pro  
 1 5 10 15  
 30 Asp Phe Ser Pro Ser Val Asp Ala Glu Ala Ile Gln Lys Ala Ile Arg  
 20 25 30  
 Gly Ile Gly Thr Asp Glu Lys Met Leu Ile Ser Ile Leu Thr Glu Arg  
 35 35 40 45  
 Ser Asn Ala Gln Arg Gln Leu Ile Val Lys Glu Tyr Gln Ala Ala Tyr  
 50 55 60  
 40 Gly Lys Glu Leu Lys Asp Asp Leu Lys Gly Asp Leu Ser Gly His Phe  
 65 70 75 80  
 Glu His Leu Met Val Ala Leu Val Thr Pro Pro Ala Val Phe Asp Ala

85

90

95

5

Lys Gln Leu Lys Lys Ser Met Lys Gly Ala Gly Thr Asn Glu Asp Ala  
 100 105 110

Leu Ile Glu Ile Leu Thr Thr Arg Thr Ser Arg Gln Met Lys Asp Ile  
 115 120 125

10

Ser Gln Ala Tyr Tyr Thr Val Tyr Lys Lys Ser Leu Gly Asp Asp Ile  
 130 135 140

Ser Ser Glu Thr Ser Gly Asp Phe Arg Lys Ala Leu Leu Thr Leu Ala  
 145 150 155 160

15

Asp Gly Arg Arg Asp Glu Ser Leu Lys Val Asp Glu His Leu Ala Lys  
 165 170 175

Gln Asp Ala Gln Ile Leu Tyr Lys Ala Gly Glu Asn Arg Trp Gly Thr  
 180 185 190

20

Asp Glu Asp Lys Phe Thr Glu Ile Leu Cys Leu Arg Ser Phe Pro Gln  
 195 200 205

25

Leu Lys Leu Thr Phe Asp Glu Tyr Arg Asn Ile Ser Gln Lys Asp Ile  
 210 215 220

Val Asp Ser Ile Lys Gly Glu Leu Ser Gly His Phe Glu Asp Leu Leu  
 225 230 235 240

30

Leu Ala Ile Val Asn Cys Val Arg Asn Thr Pro Ala Phe Leu Ala Glu  
 245 250 255

Arg Leu His Arg Ala Leu Lys Gly Ile Gly Thr Asp Glu Phe Thr Leu  
 260 265 270

35

Asn Arg Ile Met Val Ser Arg Ser Glu Ile Asp Leu Leu Asp Ile Arg  
 275 280 285

40

Thr Glu Phe Lys Lys His Tyr Gly Tyr Ser Leu Tyr Ser Ala Ile Lys  
 290 295 300

Ser Asp Thr Ser Gly Asp Tyr Glu Ile Thr Leu Leu Lys Ile Cys Gly

305

310

315

320

Gly Asp Asp

5

&lt;210&gt; 6

&lt;211&gt; 346

&lt;212&gt; PRT

&lt;213&gt; human

&lt;400&gt; 6

10

Met Ala Met Val Ser Glu Phe Leu Lys Gln Ala Trp Phe Ile Glu Asn  
 1 5 10 15

Glu Glu Gln Glu Tyr Val Gln Thr Val Lys Ser Ser Lys Gly Gly Pro  
 20 25 30

15

Gly Ser Ala Val Ser Pro Tyr Pro Thr Phe Asn Pro Ser Ser Asp Val  
 35 40 45

20

Ala Ala Leu His Lys Ala Ile Met Val Lys Gly Val Asp Glu Ala Thr  
 50 55 60

Ile Ile Asp Ile Leu Thr Lys Arg Asn Asn Ala Gln Arg Gln Gln Ile  
 65 70 75 80

25

Lys Ala Ala Tyr Leu Gln Glu Thr Gly Lys Pro Leu Asp Glu Thr Leu  
 85 90 95

Lys Lys Ala Leu Thr Gly His Leu Glu Glu Val Val Leu Ala Leu Leu  
 100 105 110

30

Lys Thr Pro Ala Gln Phe Asp Ala Asp Glu Leu Arg Ala Ala Met Lys  
 115 120 125

35

Gly Leu Gly Thr Asp Glu Asp Thr Leu Ile Glu Ile Leu Ala Ser Arg  
 130 135 140

Thr Asn Lys Glu Ile Arg Asp Ile Asn Arg Val Tyr Arg Glu Glu Leu  
 145 150 155 160

40

Lys Arg Asp Leu Ala Lys Asp Ile Thr Ser Asp Thr Ser Gly Asp Phe  
 165 170 175

	Arg	Asn	Ala	Leu	Leu	Ser	Leu	Ala	Lys	Gly	Asp	Arg	Ser	Glu	Asp	Phe
				180					185					190		
5	Gly	Val	Asn	Glu	Asp	Leu	Ala	Asp	Ser	Asp	Ala	Arg	Ala	Leu	Tyr	Glu
			195					200					205			
	Ala	Gly	Glu	Arg	Arg	Lys	Gly	Thr	Asp	Val	Asn	Val	Phe	Asn	Thr	Ile
		210					215					220				
10	Leu	Thr	Thr	Arg	Ser	Tyr	Pro	Gln	Leu	Arg	Arg	Val	Phe	Gln	Lys	Tyr
	225					230				235					240	
	Thr	Lys	Tyr	Ser	Lys	His	Asp	Met	Asn	Lys	Val	Leu	Asp	Leu	Glu	Leu
15					245					250					255	
	Lys	Gly	Asp	Ile	Glu	Lys	Cys	Leu	Thr	Ala	Ile	Val	Lys	Cys	Ala	Thr
			260					265					270			
20	Ser	Lys	Pro	Ala	Phe	Phe	Ala	Glu	Lys	Leu	His	Gln	Ala	Met	Lys	Gly
		275						280					285			
	Val	Gly	Thr	Arg	His	Lys	Ala	Leu	Ile	Arg	Ile	Met	Val	Ser	Arg	Ser
		290					295					300				
25	Glu	Ile	Asp	Met	Asn	Asp	Ile	Lys	Ala	Phe	Tyr	Gln	Lys	Met	Tyr	Gly
	305					310				315					320	
	Ile	Ser	Leu	Cys	Gln	Ala	Ile	Leu	Asp	Glu	Thr	Lys	Gly	Asp	Tyr	Glu
30				325					330					335		
	Lys	Ile	Leu	Val	Ala	Leu	Cys	Gly	Gly	Asn						
			340						345							

35

40